arXiv:1407.6233v1 [math.AP] 23 Jul 2014

A SHARP INEQUALITY FOR SOBOLEV FUNCTIONS

PEDRO M. GIRÃO

ABSTRACT. Let $N \geq 5$, $a > 0$, Ω be a smooth bounded domain in $\mathbb{R}^N$, $2^* = \frac{2N}{N-2}$, $2^\# = \frac{2(N-1)}{N-2}$ and $||u||^2 = |\nabla u|_2^2 + a|u|_2^2$. We prove there exists an $\alpha_0 > 0$ such that, for all $u \in H^1(\Omega) \setminus \{0\}$,

$$\frac{S}{2^{\frac{2}{N}}} \leq \frac{||u||^2}{|u|_{2^*}^2} \left(1 + \alpha_0 \frac{|u|_{2^\#}^{2^\#}}{||u|| \cdot |u|_{2^*}^{2^*/2}}\right).$$

This inequality implies Cherrier's inequality.

Let $N \geq 5$, $a > 0$, $\alpha \geq 0$, Ω be a smooth bounded domain in $\mathbb{R}^N$, $2^* = \frac{2N}{N-2}$ and $2^\# = \frac{2(N-1)}{N-2}$. We regard a as fixed and α as a parameter. Denote the L^p and H^1 norms of u in Ω by

$$|u|_p := (\int |u|^p)^{\frac{1}{p}} \qquad \text{and} \qquad ||u|| := \left(|\nabla u|_2^2 + a|u|_2^2\right)^{\frac{1}{2}},$$

respectively. All our integrals are over Ω. We define the functional $\delta : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$, homogeneous of degree zero, by

$$\delta(u) := \frac{|u|_{2^\#}^{2^\#}}{||u|| \cdot |u|_{2^*}^{2^*/2}}$$

and consider the system

$$\begin{cases} \left(1 + \frac{1}{2}\alpha\delta(u)\right)(-\Delta u + au) + \frac{2^\#}{2}\alpha u^{2^\#-1} = \left(1 + \frac{2^\#+1}{2}\alpha\delta(u)\right) u^{2^*-1} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases} \tag{1}_\alpha$$

We claim that the solutions of $(1)_\alpha$ correspond to critical points of the functional $\Phi_\alpha : H^1(\Omega) \setminus \{0\} \to \mathbb{R}$, defined by

$$\Phi_\alpha(u) := \left(\frac{1}{2}||u||^2 - \frac{1}{2^*}|u|_{2^*}^{2^*}\right)\left(1 + \alpha\frac{|u|_{2^\#}^{2^\#}}{||u|| \cdot |u|_{2^*}^{2^*/2}}\right)^{\frac{N}{2}} = \Phi_0(u)(1 + \alpha\delta(u))^{\frac{N}{2}}. \tag{2}$$

In fact, since

$$\Phi'_\alpha = (1 + \alpha\delta)^{\frac{N}{2}-1}\left[\Phi'_0(1 + \alpha\delta) + \frac{N}{2}\Phi_0\alpha\delta'\right]$$

and, for $\varphi \in H^1(\Omega)$,

$$\begin{aligned} \delta'(u)(\varphi) \quad = \quad & -\frac{\delta(u)}{||u||^2}\int(\nabla u \cdot \nabla\varphi + au\varphi) + 2^\#\frac{\delta(u)}{|u|_{2^\#}^{2^\#}}\int(|u|^{2^\#-2}u\varphi) \\ & -\frac{2^*}{2}\frac{\delta(u)}{|u|_{2^*}^{2^*}}\int(|u|^{2^*-2}u\varphi), \end{aligned}$$

2000 *Mathematics Subject Classification.* 46E35, 35J65.

the critical points of Φ_α satisfy

$$\begin{aligned}(-\Delta u + au)\left(1 + \tfrac{4-N}{4}\alpha\delta(u) + \tfrac{N-2}{4}\tfrac{|u|_{2^*}^{2^*}}{||u||^2}\alpha\delta(u)\right)\\ + \tfrac{2^\# N}{2}\alpha|u|^{2^\#-2}u\left(\tfrac{1}{2}\tfrac{||u||}{|u|_{2^*}^{2^*/2}} - \tfrac{1}{2^*}\tfrac{|u|_{2^*}^{2^*/2}}{||u||}\right)\\ - |u|^{2^*-2}u\left(1 + \tfrac{4-N}{4}\alpha\delta(u) + \tfrac{2^*N}{8}\tfrac{||u||^2}{|u|_{2^*}^{2^*}}\alpha\delta(u)\right) \;=\; 0\end{aligned}$$

in Ω, $\left(\frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega\right)$. However, multiplying this equation by u and integrating over Ω (i.e. differentiating (2) along the radial direction) we get $||u||^2 = |u|_{2^*}^{2^*}$. Conversely, the solutions of $(1)_\alpha$ are solutions of the previous equation multiplying $(1)_\alpha$ by u and integrating over Ω we get a quadratic equation in $||u||/|u|_{2^*}^{2^*/2}$, whose solution is $||u||/|u|_{2^*}^{2^*/2} = 1$. This proves our claim.

The functional Φ_α restricted to the Nehari manifold,

$$\mathcal{N} := \left\{u \in H^1(\Omega)\setminus\{0\} : \Phi'_\alpha(u)u = 0\right\} = \left\{u \in H^1(\Omega)\setminus\{0\} : ||u||^2 = |u|_{2^*}^{2^*}\right\},$$

is $\frac{1}{N}[\beta(1+\alpha\delta)]^{\frac{N}{2}}$, where $\beta : H^1(\Omega)\setminus\{0\} \to \mathbb{R}$ is defined by

$$\beta(u) := \frac{||u||^2}{|u|_{2^*}^2}.$$

So, we consider the functional $\Psi_\alpha : H^1(\Omega)\setminus\{0\} \to \mathbb{R}$, defined by

$$\Psi_\alpha := \beta(1+\alpha\delta).$$

A *least energy* solution of $(1)_\alpha$ is a function $u \in H^1(\Omega)\setminus\{0\}$, such that

$$\Phi_\alpha(u) = \inf_{\mathcal{N}} \Phi_\alpha = \inf_{H^1(\Omega)\setminus\{0\}} \tfrac{1}{N}(\Psi_\alpha)^{\frac{N}{2}}.$$

We are interested in proving existence and nonexistence of least energy solutions of $(1)_\alpha$. We note that every critical point of Φ_α is a critical point of Ψ_α. It is easy to check that the Nehari manifold is a natural constraint for Φ_α. So conversely, if u is a critical point of Ψ_α, then there exists a unique $t(u) > 0$, such that $t(u)u$ is a critical point of Φ_α $\left(t(u) = \left(||u||^2/|u|_{2^*}^{2^*}\right)^{\frac{N-2}{4}}\right)$.

We consider the minimization problem corresponding to

$$S_\alpha := \inf\left\{\Psi_\alpha(u) | u \in H^1(\Omega)\setminus\{0\}\right\}.$$

We recall that $S := \inf\left\{\left.\frac{|\nabla u|^2_{L^2(\mathbb{R}^N)}}{|u|^2_{L^{2^*}(\mathbb{R}^N)}}\right| u \in L^{2^*}(\mathbb{R}^N), \nabla u \in L^2(\mathbb{R}^N), u \neq 0\right\}$ is achieved by the instanton $U(x) := \left(\frac{N(N-2)}{N(N-2)+|x|^2}\right)^{\frac{N-2}{2}}$. Our main result is

Theorem 1. *There exists a positive real number*

$$\alpha_0 = \alpha_0(a,\Omega) = \min\left\{\alpha \mid S_\alpha = S/2^{\frac{2}{N}}\right\}$$

such that

(i) if $\alpha < \alpha_0$, then $(1)_\alpha$ has a least energy solution u_α;
(ii) if $\alpha > \alpha_0$, then $(1)_\alpha$ does not have a least energy solution and $\Psi_\alpha \geq S/2^{\frac{2}{N}}$. The constant $S/2^{\frac{2}{N}}$ is sharp.

Remark 1. *Obviously, α_0 is a nonincreasing function of a. By testing Ψ_α with constant functions and instantons we can prove that*

$$\alpha_0 \geq \max\left\{[S/(2|\Omega|)^{\frac{2}{N}} - a]/\sqrt{a}, C(N)\max_{\partial\Omega} H\right\}$$

where $|\Omega|$ is the Lebesgue measure of Ω, H is the mean curvature of $\partial\Omega$ and $C(N)$ is a constant that only depends on N. The least energy solutions might be constant ($a^{\frac{N-2}{4}}$) for $a \leq S/(2|\Omega|)^{\frac{2}{N}}$ if $\alpha \leq [S/(2|\Omega|)^{\frac{2}{N}} - a]/\sqrt{a}$.

Corollary 2. *For all $u \in H^1(\Omega) \setminus \{0\}$,*

$$\frac{S}{2^{\frac{2}{N}}} \leq \frac{||u||^2}{|u|_{2^*}^2}\left(1 + \alpha_0 \frac{|u|_{2^\#}^{2^\#}}{||u|| \cdot |u|_{2^*}^{2^*/2}}\right).$$

Corollary 3. *For all $u \in H^1(\Omega)$,*

$$\frac{S}{2^{\frac{2}{N}}}|u|_{2^*}^2 \leq ||u||^2 + \alpha_0||u|| \cdot |u|_2 \qquad \text{and} \qquad \frac{S}{2^{\frac{2}{N}}}|u|_{2^*}^2 \leq \left(|\nabla u|_2 + c_{a,\alpha_0}|u|_2\right)^2,$$

with $c_{a,\alpha_0} = \max\left\{\alpha_0/2, \sqrt{a + \alpha_0\sqrt{a}}\right\}$.

Proof. From Hölder's inequality $|u|_{2^\#}^{2^\#} \leq |u|_2 |u|_{2^*}^{2^*/2}$, so $\delta(u) \leq \frac{|u|_2}{||u||}$. □

Corollary 4. *(Cherrier's inequality). Let $\varepsilon > 0$. For all $u \in H^1(\Omega)$,*

$$\frac{S}{2^{\frac{2}{N}}}|u|_{2^*}^2 \leq (1+\varepsilon)||u||^2 + \frac{\alpha_0^2}{4\varepsilon}|u|_2^2 = (1+\varepsilon)|\nabla u|_2^2 + \left(\frac{\alpha_0^2}{4\varepsilon} + a\varepsilon\right)|u|_2^2.$$

Sketch of the proof of Theorem 1. – By testing Ψ_α with instantons, $S_\alpha \leq \frac{S}{2^{\frac{2}{N}}}$, for all $\alpha \geq 0$. We claim that if $S_\alpha < \frac{S}{2^{\frac{2}{N}}}$, then S_α is achieved. This is a consequence of the concentration-compactness principle. A minimizing sequence u_k with $|u_k|_{2^*} = 1$ is bounded and we can assume $u_k \rightharpoonup u$ in $H^1(\Omega)$, $\lim_{k\to\infty}|\nabla u_k|_2^2 = |\nabla u|_2^2 + ||\mu||$ and $\lim_{k\to\infty}|u_k|_{2^*}^{2^*} = |u|_{2^*}^{2^*} + ||\nu|| = 1$, where $\frac{S}{2^{\frac{2}{N}}}||\nu||^{\frac{2}{2^*}} \leq ||\mu||$ (we remark that this inequality follows from Cherrier's inequality). We can write $\beta(u)\delta(u) = \gamma(u)\sqrt{\beta(u)}$ for $\gamma(u) = |u|_{2^\#}^{2^\#}/|u|_{2^*}^{2^\#}$. The key step in the proof of the claim is the following observation. Define f and $g : [0,1] \to \mathbb{R}$, by

$$\begin{aligned} f(x) &:= \beta x^{\frac{2}{2^*}} + \frac{S}{2^{\frac{2}{N}}}(1-x)^{\frac{2}{2^*}} + \alpha\gamma x^{\frac{2^\#}{2^*}}\sqrt{\beta x^{\frac{2}{2^*}} + \frac{S}{2^{\frac{2}{N}}}(1-x)^{\frac{2}{2^*}}} \\ &\geq \beta x + \frac{S}{2^{\frac{2}{N}}}(1-x) + \alpha\gamma x\sqrt{\beta x + \frac{S}{2^{\frac{2}{N}}}(1-x)} \quad =: \quad g(x). \end{aligned}$$

Suppose $\min f < \frac{S}{2^{\frac{2}{N}}}$. It follows that $\beta < \frac{S}{2^{\frac{2}{N}}}$, and this in turn implies that g is concave. Since f and g coincide at 0 and 1, the minimum of f occurs at 1. This proves the claim.

A similar argument shows $\alpha \mapsto S_\alpha$ is continuous. In particular, the supremum $\alpha_0 := \sup\{\alpha | S_\alpha < S/2^{\frac{2}{N}}\}$ is either $+\infty$ or a maximum. The map $\alpha \mapsto S_\alpha$ is strictly increasing on $[0, \alpha_0]$. If $\alpha \in [0, \alpha_0[$, then $(1)_\alpha$ has a least energy solution. If $\alpha \in]\alpha_0, +\infty[$, then $(1)_\alpha$ does not have a least energy solution. It remains to prove that α_0 is finite.

Suppose $\alpha_0 = +\infty$. Choose $\alpha_k \to +\infty$ as $k \to +\infty$ and let u_k be minimizer for S_{α_k} satisfying $(1)_{\alpha_k}$. It is easy to prove that $\lim_{\alpha\to\infty} S_\alpha = S/2^{\frac{2}{N}}$, $M_k := \max_{\bar\Omega} u_k = u_k(P_k) \to +\infty$, $\lim_{k\to\infty} |\nabla u_k|_2^2 = \lim_{k\to\infty} |u_k|_{2^*}^{2^*} = S^{\frac{N}{2}}/2$ and $\alpha_k \delta(u_k) \to 0$. We can apply the Gidas-Spruck blow up technique to $(1)_{\alpha_k}$ because $\alpha_k\delta(u_k) \to 0$. Define $\varepsilon_k := M_k^{-2/(N-2)}$ and $U_{\varepsilon,y} := \varepsilon^{-\frac{N-2}{2}} U\left(\frac{x-y}{\varepsilon}\right)$. We can prove that $\lim_{k\to\infty} \alpha_k\varepsilon_k = 0$, $\lim_{k\to\infty} |\nabla u_k - \nabla U_{\varepsilon_k,P_k}|_2 = 0$ and $P_k \in \partial\Omega$, for large k.

At this point, using the ideas of [2], we follow the argument in [7], which applies with no modification. We show $\Psi_{\alpha_k}(u_k) > S/2^{\frac{2}{N}}$, for large k. This is impossible. Therefore α_0 is finite. □

Remark 2. *The functional behavior behind this inequality is also present, for example, in the Dirichlet problem for* $-\Delta u - au + \alpha u^{1/3} = u^{7/3}$ *in* $\Omega \subset \mathbb{R}^5$, *with* $0 < a < \lambda_1\left(-\Delta, H_0^1(\Omega)\right)$. *In this case* $s := [t(u)]^{2/3}$ *is the solution of a cubic equation,* $(|\nabla u|_2^2 - a|u|_2^2)s + \alpha|u|_{4/3}^{4/3} - |u|_{10/3}^{10/3}s^3 = 0$.

References

[1] Adimurthi, Mancini G., The Neumann problem for elliptic equations with critical nonlinearity, A tribute in honour of G. Prodi, Scuola Norm. Sup. Pisa (1991) 9-25.

[2] Adimurthi, Pacella F., Yadava S.L., Interaction between the geometry of the boundary and positive solutions of a semilinear Neumann problem with critical nonlinearity, J. Funct. Anal. 113 (1993) 318-350.

[3] Brézis H., Lieb E., Sobolev inequalities with remainder terms, J. Funct. Anal. 62 (1985) 73-86.

[4] Brézis H., Nirenberg L., Positive Solutions of Nonlinear Elliptic Equations Involving Critical Sobolev Exponents, Comm. on Pure and Appl. Math., Vol. 36 (1983) 437-477.

[5] Chabrowski J., Willem M., Least energy solutions for the critical Neumann problem with a weight, to appear in Calc. Var.

[6] Cherrier P., Problémes de Neumann nonlinéaires sur des variétés Riemannienes, J. Funct. Anal. 57 (1984) 154-207.

[7] Costa D.G., Girão P.M., Existence and nonexistence of least energy solutions of the Neumann problem for a semilinear elliptic equation with critical Sobolev exponent and a critical lower-order perturbation, to appear.

[8] Lions P.L., The concentration-compactness principle in the calculus of variations, The limit case, Revista Math. Iberoamericana 1 No. 1 and No. 2 (1985) 145-201 and 45-120.

[9] Wang X.J., Neumann problems of semilinear elliptic equations involving critical Sobolev exponents, J. Diff. Eq. 93 (1991) 283-310.

[10] Wang Z.Q., Existence and nonexistence of G-least energy solutions for a nonlinear Neumann problem with critical exponent in symmetric domains, Calc. Var. 8 (1999) 109-122.

[11] Zhu M., Sharp Sobolev inequalities with interior norms, Calc. Var. 8 (1999) 27-43.

Mathematics Department, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisbon, Portugal

E-mail address: pgirao@math.ist.utl.pt